901 45th Street, Kimmel Building West Palm Beach, FL 33407

> P: 561-844-5255 F: 561-844-5945

Ash Itinger, Admin Ext 246 www.cartilagerepaircenter.org www.paleyinstitute.org

Tom Minas, MD, MS

HTO with Autologous Bone Graft and Weight-bearing Condyle PT Protocol Stage 2- Transitional Phase (7+ weeks)

PRIMARY GOALS

DO NOT OVERLOAD GRAFT

INCREASE TIBIAL-FEMORAL AND PATELLA-FEMORAL MOBILITY RESTORE QUADRICEPS CONTROL

Weeks 7 & 8

Weeks 9 & 10

Weeks 11 & 12

PWB 1/3 Body Weight
PWB 2/3 Body Weight
FWBAT with crutches

•Knee immobilizer with ambulation until good quad control/independent SLR

Week 13+ Crutch, Cane, or no device as tolerated

*Progress per guidelines above as pain allows.

Continue full AROM and gentle PROM exercises

•CPM may be discontinued.

THEREX •Low weight (max 10-20 lbs) open-chain leg extension and curl

•Use bathroom scale to progress as follows:

•Stationary bicycle with gradual increased tension per level of comfort.

•Continue quad sets, SLR in brace, leg curl and heel slides

*Strengthen quadriceps, hamstrings, and hip abductors/extensors using ankle weights

and/or elastic band resistance through full ROM as tolerated

•Gentle closed-chain terminal knee extension 0-40 degrees (TKE) permitted starting at 9-

10 weeks as tolerated per weight bearing restriction

THERAPY •Continue gentle multi-directional patella mobilization as needed

•Whirlpool or pool therapy as available to enhance ROM and quadriceps/hamstring muscle

control

•E-stim for VMO/quadriceps muscle re-education/biofeedback as needed

•Gentle massage/deep friction to hamstring insertions, suprapatellar quadriceps,

medial/lateral gutters, and infrapatellar regions

COMMENTS • Activity level should be modified if increased pain, catching, or swelling occurs

•No progression of this protocol until cleared by MD at 12 weeks post-op

For Outpatient Physical Therapist:

2-3x week / 6 weeks

BRACE

GAIT

ROM

